

Position Statement

SUMMARY

It is the position of the National Association of School Nurses (NASN) that the management of head lice (*Pediculus humanus capitis*) in the school setting should not disrupt the educational process. Leadership provided by the registered professional school nurse (hereinafter referred to as the school nurse) can impact reduction of the stigma associated with head lice by providing accurate health education including anticipatory guidance to the school community and implementing evidence-based strategies for the management of head lice in schools. Evidence-based strategies include abandoning “no-nit” school policies, allowing children to remain in class and participate in school-sponsored activities when live lice or nits (the eggs of head lice) are found on their heads, notifying parents/caregivers at the end of the school day when findings indicate the presence of a head lice infestation, and educating parents/caregivers about evidence-based treatment options.

BACKGROUND

In the United States, head lice infestations are most common among preschool and elementary school-age children and their household members regardless of socioeconomic status and hygienic living conditions (Centers for Disease Control and Prevention [CDC], 2013a). According to research head lice infestations predominantly affect the age group of 3-11 years (Frankowski & Bocchini, 2010), with an estimated 6 million to 12 million cases annually (CDC, 2013a). A 2004 study estimated annual direct and indirect costs associated with head lice infestations and recent treatment costs at \$1 billion (Hansen & O’Hayer, 2004). “No-nit” policies that require a child to be free of nits before he or she can return to school lack evidence of being effective, result in unnecessary absenteeism, and may violate affected children’s civil liberties (Pontius, 2014; CDC, 2013a). Unnecessary absenteeism leads to missed learning opportunities for the student and potentially lost family wages due to loss of parent/guardian workdays (Pontius, 2014).

Head lice are not known to cause disease; however, secondary bacterial infection of the skin resulting from contaminated scratching and related lesions can occur. Research has shown that the survival of head lice when not on the head is usually less than one day, and the eggs can only hatch when incubated by body heat found near the scalp (Devore et al., 2015; CDC, 2013c). Transmission occurs primarily through head-to-head contact and infrequently through indirect contact with shared personal belongings.

Even with this knowledge, the presence of head lice can negatively affect families and schools. For the student and family there can be significant social stigma and caregiver strain (Gordon, 2007). For the school, when evidence-based policies and intervention strategies are not in place, head lice can significantly disrupt the education process (CDC, 2013c; Pontius, 2014).

In the past, many schools with “no nit” policies expended innumerable hours and resources in attempts to eradicate head lice infestations. Studies have shown that control measures such as, mass screenings for nits, have not been shown to have a significant effect on the incidence of head lice in a school community, nor have they shown to be cost-effective (Devore et al., 2015; Meinking & Taplin, 2011; CDC, 2013a). Communication between school personnel and parents/caregivers highlighting cases of head lice (e.g., “head lice outbreak letters”) has been shown to increase community anxiety, increase social stigma causing embarrassment of affected infested students, and puts students’ rights to confidentiality at risk (Gordon, 2007; Pontius, 2014).

Head lice treatment success is variable, adding to confusion and frustration among students, families, and members of the school community. Some children develop persistent head lice, which requires-concentrated efforts to address treatment as well as the stress experienced by the child and family (Gordon, 2007). Head lice in some communities have developed resistance to common over-the-counter treatments, resulting in the need for a more individualized approach to management by a healthcare provider (Yoon et al., 2014; Meinking et al., 2002;

Devore et al., 2015). Treatment failures can also result from initial misdiagnosis, non-adherence to a treatment protocol, a new infestation acquired after treatment, or the lack of use of an ovicidal product (Devore et al., 2015; Pontius; 2014; Pollack, Kiszewski, & Spielman, 2000; CDC, 2013b).

RATIONALE

Evidence-based strategies for the management of head lice in the school setting can reduce the incidence of infestations, the social stigma and caregiver strain experienced by students and families, and the negative impact on students' education. The school nurse can provide leadership within the school community to effectively manage head lice by:

- Attaining knowledge and competency that reflect current evidence-based school nursing practice related to the management of head lice (American Nurses Association & National Association of School Nurses [ANA & NASN], 2011).
- Providing accurate health education to the school community focused on dispelling common myths about head lice (e.g., incidence, life cycle of the head louse, mode of transmission, importance of regular surveillance at home, recommended evidence-based treatment options, care of the environment) (ANA & NASN, 2011; Pontius, 2014).
- Advocating and providing rationale for the elimination of mass school screenings for head lice (Devore et al., 2015; CDC, 2013a).
- Educating families about how to assess their children for suspected head lice (Devore et al., 2015).
- Providing privacy when conducting student health assessment for suspected or reported cases of head lice (ANA & NASN, 2011).
- Returning affected students to class or other school sponsored activities with instruction to avoid head-to-head contact (Pontius, 2014). If live lice or nits are found,
 - Eliminating classroom-wide or school-wide family head lice notification.
 - Notifying parents/caregivers at the end of the school day to teach about evidence-based treatment options and steps to follow.
- Advocating for and providing rationale for the abandonment of "no-nit" school policies that require a child to be free of nits before he or she can return to school (Devore et al., 2015; Pontius, 2014).
- Educating parents/caregivers about the chosen evidence-based treatment option, the importance of adherence with the treatment protocol, and the importance of reassessment for recurrence (Devore et al., 2015; Pontius, 2014).

CONCLUSION

The school nurse is the health professional who provides leadership for the school community to implement evidence-based strategies for the management of head lice in the school setting. The role of the school nurse includes the following (Pontius, 2014; Devore et al., 2015; CDC, 2013a):

- Provide accurate health education to the school community about the etiology, transmission, assessment, and treatment of head lice;
- Advocate for school policy that is more caring and less exclusionary (i.e., elimination of the "no-nit" school policies);
- Implement intervention strategies that are student-centered;
- Support the current treatment recommendation of the American Academy of Pediatrics and CDC; and
- Participate in research that evaluates the effectiveness of head lice policies and educational programs.

It is unlikely that all head lice infestations can be prevented. Parents/caregivers will benefit from receiving support from the school nurse about the importance of regular surveillance at home, choosing and adhering to the protocols of evidence-based treatment recommendations, and educating to dispel head lice myths. The education mission of schools will be supported by implementing evidence-based policies and strategies under the guidance of the school nurse. The burden of unnecessary absenteeism to the students, families, and communities far outweighs the perceived risks associated with head lice.

REFERENCES

- American Nurses Association & National Association of School Nurses. (2011). *School nursing: Scope and standards of practice* (2nd ed.). Silver Spring, MD: Nursesbooks.org.
- Centers for Disease Control and Prevention. (2013a). *Head lice information for schools*. Retrieved from <http://www.cdc.gov/parasites/lice/head/schools.html>
- Centers for Disease Control and Prevention. (2013b). *Parasites – lice – head lice - treatment*. Retrieved from <http://www.cdc.gov/parasites/lice/head/treatment.html>
- Centers for Disease Control and Prevention. (2013c). *Biology*. Retrieved from www.cdc.gov/parasites/lice/head/biology.html
- Devore, C., Schutze, G., & The American Academy of Pediatrics' Council On School Health and Committee on Infectious Diseases. (2015). Head lice, *Pediatrics*, 135(5), e1355-e1365. Doi:10.1542/eds.2015-0746
- Gordon, S. (2007). Shared vulnerability: A theory of caring for children with persistent head lice. *The Journal of School Nursing*, 23(5), 283-294.
- Frankowski, B.L., & Bocchini, J.A. (2010). Clinical report-head lice. *Pediatrics*, 126(2) 392-403. Retrieved from <http://pediatrics.aappublications.org/cgi/content/abstract/126/2/392>
- Hansen, R., & O'Haver, J. (2004). Economic considerations associated with *Pediculus humanus capitis* infestation. *Clinical Pediatrics*, 43(6), 523-527.
- Meinking, T., & Taplin, D. (2011). Infestations. In L. Schachner, & R. Hansen (Eds.). *Pediatric dermatology* (pp. 1525-1583). Philadelphia, PA: Mosby Elsevier.
- Meinking, T., Serrano, L., Hard, B., Entzel, P. Lemard, G., & Villar, M. (2002). Comparative in vitro pediculicidal efficacy of treatments in a resistant head lice population in the United States. *Archives of Dermatology*, 138(2), 220-224.
- Pollack, R., Kiszewski, A., & Spielman, A. (2000). Over diagnosis and consequent mismanagement of head louse infestations in North America. *Pediatric Infectious Disease Journal*, 19(8), 689-693.
- Pontius, D. (2014). Demystifying pediculosis: School nurses taking the lead. *Pediatric Nursing*, 40(5), 226-235. Retrieved from <https://www.pediatricnursing.net/ce/2016/article4005226235.pdf>
- Yoon, K., Previte, D., Hodgdon, H., Poole, B, Kwon, D., El-Ghar, G., Clark, M. (2014). Knockdown resistance allele frequencies in North American head louse (Anoplura: Pediculidae) populations. *Journal of Medical Entomology*, 51(2), 450-457. doi: <http://dx.doi.org/10.1603/ME13139>

Acknowledgment of Authors:

Suzanne Smith, BSN, RN, NCSN
Nichole Bobo, MSN, RN
Kathy M. Strasser, MS, RN, NCSN
Kathey M. Haynie, MSN, RN

Adopted: 1999

Revised: 2004, January 2011, January 2016


This document replaces the position statement titled *Pediculosis Management in the School Setting, 2011*.

Suggested citation: National Association of School Nurses. (2016). *Head lice management in the school setting* (Position Statement). Silver Spring, MD: Author.

All position statements from the National Association of School Nurses will automatically expire five years after publication unless reaffirmed, revised, or retired at or before that time.

www.nasn.org
National Association of School Nurses
1100 Wayne Avenue, Suite 925
Silver Spring, MD 20910
1-240-821-1130

Figure 2.
Lifecycle of the Louse



Source: CDC, 2013c.

Figure 3.
Viable Nit




Source: CDC Public Health Image Library, 2014.

2013a). Considering that nits do not move after being laid, that nits hatch in approximately seven days, and that hair grows and average of 13 mm (Caucasian) to 10 mm (African American) per month (Loussouarn, Rawadi, & Genain, 2005), and generously doubling the viability estimate (as it can be longer in warmer climates (Meinking & Taplin, 2011), most experts concur that nits found farther than one-fourth to one-half inches (6 to 12mm) from the scalp are non-viable (Frankowski & Bocchini, 2010; Mumcuoglu et al., 2007; National Association of School Nurses (NASN), 2011; Pollack et al., 2000). Lice are more common among girls, which may be due to longer hair hiding the infestation or a greater likelihood of playing with heads closer together (Burgess et al., 2003; Frankowski & Bocchini, 2010). By the time a case of lice is found, the child has generally had them for a month or more (Frankowski & Bocchini).

Myth #6. No-Nit Policies Reduce the Transmission Of Head Lice in Schools

Over the past decade, there have been important recommended changes in the management of head lice in schools: *No student should ever miss school time because of head lice, and no-nit policies should be eliminated* (CDC, 2013a; Frankowski & Bocchini, 2010; Mumcuoglu et al., 2007; NASN, 2011). Many schools have traditionally had “no-nit” policies, which require the removal of all nits, viable or not, before a child returns to school. To school personnel unfamiliar with the life cycle of the louse, school exclusion for an infestation or for the evi-

Figure 4.
Comparison of Egg, Nymph, and Adult



Source: CDC Public Health Image Library, 2014.

dence of nits seems logical. Their rationale may be that by excluding those who have head lice, others will not catch it. However, there is no evidence that these policies reduce the transmission of head lice in schools. Moreover, there is significant evidence they increase absenteeism, shame, stigma, and unnecessary treatment.

The Cost of “No-Nit” Policies

It has been estimated that school children with lice lose an average of four days of school per year in schools where “no-nit” policies are enforced (Gordon, 2009). The loss of the opportunity for learning, funding for schools, and parent/guardian work days result. Our nation’s schools lose between \$280 to \$325 million in annual funding, and families lose up to \$2,720 in wages per active infestation (Gordon, 2009). One study found that while the presence of more than five nits closer than one-quarter inch from the scalp *was* a risk factor for the development of an active infestation, most of these children did not actually become infested (Williams, Reichert, MacKenzie, Hightower, & Blake, 2001). Just the presence of nits does not indicate the presence of an active case of lice, especially if the nits are more than approximately one half inch (1 cm) from the scalp.

Even viable nits do not transmit lice. Eggs cannot be transmitted from one head to another, nor can they fall off the hair shaft. Even if hair with a viable nit falls off, it will not hatch at temperatures lower than the human head (Meinking & Taplin, 2011). Should environmental temperatures stay warm enough that the nymph actually hatches *off* the head, as an obligate ectoparasite and blood feeder, it must find a human host or rapidly succumb within 24 to 48 hours (Frankowski & Bocchini, 2010). There is no medical need to eliminate empty egg cases, but removal for esthetic reasons may reduce stigma (Burgess et al., 2003; Gordon, 2007). Additionally, the evidence shows both lay and health care personnel, even school nurses (who generally spend the most time of any health care professional assessing for lice) actually identify hair casts, hair product debris, dandruff, and other items found in the hair erroneously as lice or nits equally as often as they do so correctly (Pollack et al., 2000).

Exclusion for Live Lice

It also is no longer recommended to exclude children *immediately* for live lice or viable nits, but rather, to wait to notify parents/guardians at the end of the day. In most situations, the child has probably had lice for a month or more and possesses little risk to others (CDC, 2013a; Frankowski & Bocchini, 2010; Mumcuoglu et al., 2007). Any exposure to his or her classmates has already occurred, and immediate exclusion provides no further prevention. No exclusion from any activities is necessary, including riding the school bus or participating in sports.

Shame and Stigma

When a child is called to the nurse’s office and does not return, and then a note goes home at the end of the day to check children for possible lice, it only takes a few questions from parents/guardians to determine which of their child’s classmates has lice. For the child with persistent lice, the shame and stigma can be devastating not only to the child, but to the family as well. Children may be told they cannot play, sit by, or even be friends with the child who had/has lice. Head lice are not a health threat. Unlike body lice, head lice cause no known disease other than the occasional topical infection from persistent scratching, yet can cause a child to be socially ostracized (Gordon, 2007)

The Call for the Discontinuance

The following groups all call for elimination of “no-nit” policies:

- American Pediatrics Association (AAP) (Frankowski & Bocchini, 2010).
- Centers for Disease Control and Prevention (CDC, 2013a).
- International Guidelines for the Treatment of Pediculosis (Mumcuoglu et al., 2007).
- National Association of School Nurses (NASN, 2011).

According to D. Taplin, “If no nit policies were that effective, why do we still have so many head lice?” (Burgess et al., 2003, p. 11). However, eliminating “no nit” policies does not mean eliminating a need to treat the infestation. Whether by chemical or mechanical means, treatment to eliminate the head lice remains a high priority. Although not dangerous, infestation may be uncomfortable and should be managed.

Myth #7. Schools Are a Common Place for Lice Transmission

Surprisingly, schools rarely provide an opportunity for close head-to-head contact, except for very young children, such as preschool and kindergarten students. For that reason, schools are rarely a source for lice transmission. Head lice are most often a community health issue brought into the school setting. Speare, Thomas, and Cahill (2002) found that while 14,000 live lice were found on the heads of 466 children, no lice were found on the carpets of 118 classrooms. Hootman (2002) mapped classrooms of infested students, and found all students in the same classroom with lice shared time together outside of school with relatives, or household members, or had participated in a recent sleepover. Clothing stored next to each other, classroom headphones, riding on the bus together, and playing on the playground or in sports are also *not* sources of transmission (Burgess et al., 2003). The evidence indicates 1% to 10% of U.S. children (in kindergarten to fourth grade) have an infestation of head lice at any one time (Pollock et al., 2000). It is estimated that 10% of those may actually be transmitted in school. It does not make sense to exclude children when the likelihood of transmission in school is only 1%, far less than the common cold.

Schools often see a spike in cases after a school break, such as the beginning of the school year, after Christmas, and again after spring break. This is often falsely attributed to a return to the school environment, but is actually due to *being in the community* for an extended period of time (Gordon, 2007). These break times are commonly when children have sleepovers, go to camp, or visit relatives. They then return to school, and the teacher or school nurse who is familiar with the symptoms identifies the infestation. The school, rather than being the proximate cause of infestation, is the location of its identification.

Current recommendations include notifying parents at the *end* of the school day and providing education on the proper treatment. The child should be checked again the next school day. Should the parents be unable to provide necessary follow

through, further follow up, which may include financial assistance with pediculicides, additional education and how to check and comb out lice and viable eggs, referral for prescription treatment, or as a rare, very last resort, exclusion for non-compliance, may be appropriate (Frankowski & Bocchini, 2010; Pontius, 2011). A child should never lose a day at school because of lice.

Myth #8. Classroom Checks Can Limit Spread Of Head Lice in Schools

It is the position of NASN, the CDC, and AAP that school screenings, either routine or after an identified classroom case, are not productive, cost-effective, or merited, and are wasteful of education time (CDC, 2013a, Frankowski & Bocchini, 2010; NASN, 2011). School screenings are not an accurate way of assessing or predicting which children are or will become infested, and such screenings have not been proven to have a significant effect on the incidence of head lice in a school (Frankowski & Bocchini, 2010; Meinking & Taplin, 2011). One study found that misdiagnosis is so common that non-infested children were excluded from school *more* often than actually infested children (Pollack et al., 2000). Anecdotally, prior to the elimination of “no-nit” policies and classroom screening in her district, the author had conducted classroom screens whenever one student was found to have head lice. During eight years of such screenings, no further cases of lice were ever found that could not be attributed to close contact outside of school.

Screenings also have significant potential to violate the children’s privacy. In schools, parents or guardians have a right to control access to their child’s body. This could be violated by routinely screening students without parent/guardian permission. The National Pediculosis Association, a lay pediculosis interest group, continues to recommend the strict adherence to “no-nit” policies via nit combing and routine screenings as a way to eliminate the need any pediculicides. There is no published, reviewed evidence to support these claims (R. Pollack, personal communication, May 21, 2014).

It is prudent, however, to check close contacts of a child found to have

head lice (Frankowski & Bocchini, 2010). A close contact includes all members of the household; those who have recently spent the night; family members who travel between households in blended families; children who spend large amounts of time outside of school with each other, such as day care, camp, or at babysitters; and preschool and kindergarten children who both sit near each other and play often together.

Myth #9. Letters to Parents Or Guardians When a Case Is Identified at School Are A Good Way to Control The Spread of Head Lice

There is no evidence to support the claim that letters sent home prevent head lice transmission, and they may, in fact, be a violation of privacy and confidentiality (American School Health Association [ASHA], 2000; Frankowski & Bocchini, 2010). There is no known method to *prevent* lice (other than by shaving the scalp hair). Sending home a letter may, as it should, cause parents to check their students to see if they are currently infested. However, this may also create a false sense of security because parents may believe their child is lice-free. There may be undetected, unhatched viable nits or one pregnant louse in the hair, or the child may spend the next night with friend who unknowingly has an infestation. Some parents/guardians will treat prophylactically, causing unnecessary use of pediculicides or time-consuming combing and environmental cleaning. Sending letters home often results in panic and emotional distress among caregivers. Letters home not only provoke a crisis situation and unjustified panic, but they perpetuate the myth that lice are transmitted in schools (Mumcuoglu et al., 2007). However, some schools continue to send alert letters because while they may understand head lice are not a public health risk, they are concerned about a public relations dilemma and community backlash (Frankowski & Bocchini, 2010).

Confidentiality Violations

Parents or guardians often insist they have a right to know when a case of head lice is discovered in a classroom. Parents have also insisted they have a right to know when a child has

HIV or other communicable condition in school. However, they do *not* have a legal right to such information. Although no school would send home a letter with a specific child’s name in it, families can easily determine which child is suspected to have lice. This right to confidentiality in schools is protected by the Family Educational Rights and Privacy Act (FERPA), and by state and national ethical health care and education standards.

The Family Educational Rights and Privacy Act

FERPA requires that medical and educational records cannot, without parental/guardian consent, be released to others without a legitimate educational interest. This is regardless of whether the information is written, oral, or electronic (ASHA, 2000; Bergren, 2001). Even without disclosing the actual name, if another person can easily determine the identity of a child, then student privacy and confidentiality has been violated. For example, if the nurse were discussing an issue at school about a child in a wheelchair and only one child is in a wheelchair at school, enough information has been provided to identify that child and breach his or her privacy, without ever mentioning a name. In a pediculosis situation, if after parents receive a lice alert letter they ask their own child who went home from school today and their child knows the answer, a similar breach has occurred.

National Ethical Standards And State Laws

Both the professions of education and nursing have developed codes of ethics that stipulate not disclosing information about students obtained within the course of professional service. For example, provision #3 in the American Nurses Association (ANA) Code of Ethics states “the nurse promotes, advocates for and strives to protect the health, safety and rights of a patient, which includes both privacy and confidentiality” (ANA, 2001, p. 6). Failure to uphold national professional standards can leave the nurse open to charges of malpractice.

Harm vs. Duty to Warn

According to ASHA (2000), when contemplating a disclosure of confidential health information even if by default, two ethical criteria must be met. The criteria and related consider-

ations as they pertain to head lice include:

Ethical Criteria 1: *Do no harm.* Consider the following:

- *Can the truth of the information be confirmed?* The literature reveals how frequently health care professionals error in the identification of head lice. True confirmation must be made by microscopy.
- *How much will the individual and his or her family's privacy be violated by this disclosure? Will it harm them?* Stigma, embarrassment, and social humiliation are common for those with head lice.
- *Will a decision to disclose do MORE harm than good to the individual with head lice?* Head lice are annoying, but they are not life threatening. Stigma, however, can change lives forever.

Ethical Criteria 2: *Duty to warn.* Consider the following:

- These circumstances are limited to very few situations when the potential for harm is high.
- Examples include child abuse, self-injury, or possible life-threatening or serious harm to another person.

Because head lice are not dangerous and do not cause disease, an infestation does not rise to the level of "duty to warn." Further, because disclosure, however inadvertent, may cause great harm to the family with head lice through social stigma, the right to privacy and confidentiality must be upheld for the student and family experiencing lice. Parents/guardians are better served rather than receiving alert letters, receiving regularly scheduled head lice information letters several times during the year (most appropriately at the times when children are returning from the community after school breaks). Suggested contents include reminders to regularly check their children's hair (weekly for elementary age students) for any evidence of head lice, how to check, and treatment instructions (Gordon, 2009).

Myth #10. If One Member Of the Household Has Lice, Everyone Should Be Treated

Treatment should be initiated only when there is clear evidence of head lice. When lice are identified in

one family member, all household members and close contacts should be examined. Treat only those contacts that actually have crawling lice or viable eggs. Prophylactic treatment is unnecessary and time-consuming, and exposes persons to medications unnecessarily. All persons with head lice should be treated at the same time, otherwise they could re-infest each other (CDC, 2013a).

Myth #11. Pediculicides Are Dangerous and Should Be Avoided

Treatment choices for lice should be based on any local patterns of resistance, ease of use, and cost. Figure 5 describes a suggested treatment regimen. If the hair is fine and untangled, and the caregiver is motivated, using a fine-toothed comb and methodically combing through all hair on the head, both to examine the head for live lice, and to remove the viable nits and the lice, can be effective. This process must be repeated every few days for at least two weeks, to eliminate each new louse as it hatches. Because this is so time-consuming, and most families want the problem solved immediately, use of Federal Drug Administration (FDA)-approved pediculicides, which are safe when used as directed, can be used as an adjunct or to replace combing (Burgess et al., 2003; CDC, 2013a). The safety and effectiveness of home or "natural" remedies, such as olive oil, tea-tree oil, lavender oil, or mayonnaise, are not regulated by the FDA and have not been shown to be effective in any known double-blind studies, and are therefore not recommended (Frankowski & Bocchini, 2010)

Pediculicides

Over-the-counter (OTC) preparations or permethrin (e.g., Nix[®]) and pyrethrins (e.g., RID[®], Clear[®], Pronto[®]) remain as the first line choice for pediculicidal treatment. Even in light of some developing resistance to pyrethrins, they remain very effective. They are inexpensive and have extremely low toxicity. Pyrethrins are a natural chrysanthemum extract, and permethrin is a synthetic pyrethroid. Both are neurotoxic to lice (Frankowski & Bocchini, 2010).


Evidence has shown that many families rely on peers, and increasingly, the Internet, for their treatment

information. In addition, families will self-treat an average of five times before seeking the help of a health care professional (Gordon, 2009). The cost of seeing a health care provider, plus the additional cost of a prescription medication and the stigma associated with head lice, are often the reasons many families will initially seek alternative information and treatments. Experts recommend referral to a health care provider after two treatment failures (Burgess et al., 2003). According to the treatment protocol recommended by the American Academy of Pediatrics (2012), "For treatment failures not attributable to improper use of an over-the counter pediculicide, malathion, benzyl alcohol lotion, or spinosad suspension should be used" (p. 4). Several prescription medications exist, with three new medications receiving FDA approval in the last five years (see Table 1). Each of these pediculicides must be used according to the directions. Some require a second treatment to ensure eradication of newly hatched nymphs. Others require an extended time on the head. The school nurse can help assure treatment success by making sure explanations of options and instructions for use are clear and understood by the caregiver. Finally, there is some evidence that a device that uses hot air to desiccate the insects may be effective (Frankowski & Bocchini, 2010). However, the device is expensive and not readily available.

Lice Removal Services

Delousing/nit picking services have expanded across the county in the last decade. These primarily for-profit businesses espouse to provide relief for families who are either unable or unwilling to do mechanical lice and egg removal, and/or prefer not to use pediculicides. However, unlike barbers and beauticians who also manipulate hair, such businesses are unregulated. Workers are not health care professionals and are generally trained by the establishment (Pollack, 2012). Noted lice expert Richard Pollack (personal communication, May 21, 2014) is unconvinced of the need for such services, and believes when the nature of the business is to sell head lice control services and proprietary supplies, it is not surprising to see these salons aggressively arguing in favor of no-nit policies, and perhaps suggesting that "out-

Figure 5.
Suggested Scheme for Head Louse Infestations



Source: © 2010 IdentifyUS, LLC. Used with permission. Retrieved from <https://identify.us.com/idmybug/head-lice/head-lice-documents/lice-mgmt-chart-home.pdf>

Table 1.
Medications

	Age	Notes
Over-the-Counter Medications		
Permethrin lotion, 1%	2 months and older	<ul style="list-style-type: none"> • Kills live lice but not unhatched eggs. • A second treatment often is necessary on day 9 to kill any newly hatched lice before they can produce new eggs.
Pyrethrins	2 years and older	<ul style="list-style-type: none"> • Generally should not be used by persons who are allergic to chrysanthemums or ragweed. • A second treatment is recommended 9 to 10 days after the first treatment to kill any newly hatched lice before they can produce new eggs.
Prescription Medications		
Benzyl alcohol lotion (0.5%)	6 months and older	<ul style="list-style-type: none"> • Kills lice but not eggs. • A second treatment is needed 7 days after the first treatment to kill any newly hatched lice before they can produce new eggs.
Ivermectin lotion, 0.5%	6 months and older	<ul style="list-style-type: none"> • Kills live lice and appears to prevent nymphs (newly hatched lice) from surviving. • It is effective in most patients when given as a single application on dry hair without nit combing. • It should not be used for retreatment without talking to a health care provider.
Malathion lotion, 0.5%	6 years and older	<ul style="list-style-type: none"> • Kills live lice and some lice eggs. • A second treatment is recommended if live lice still are present 7 to 9 days after treatment.
Spinosad 0.9% topical suspension	4 years and older	<ul style="list-style-type: none"> • Kills live lice and unhatched eggs. • Retreatment usually not needed and should be given only if live (crawling) lice are seen 7 days after first treatment.

Source: Adapted from CDC, 2013b.

breaks” or “epidemics” of head lice are occurring. The cost can be quite expensive, with a session costing several hundred dollars or more. There is no evidence to support recommending these services.

Myth #12. Head Lice Are Becoming Increasingly Resistant to Pediculicides

Several studies have reported some increase in local resistance to OTC pediculicides in the last 20 years, receiving prominent lay press coverage. However, the prevalence of actual resistance is unknown because clinical trials have used different inclusion criteria, resulting in different conclusions (Frankowski & Bocchini, 2010; Pollack et al., 2000). Most recently, Yoon et al. (2014) found by DNA typ-

ing of lice in several locations in the U.S. and Canada, the rate of T1 mutation (the gene mutation most responsible for permethrin resistance) varied between 84.4% and 99%. While this suggests increasing resistance to permethrin and pyrethrins-based pediculicides, one must be careful in extrapolating these results because their study examined a small number of lice from only 12 U.S. states and studied the potential for resistance, rather than a clinical measurement of actual resistance. What matters most is the degree to which head lice have become resistant in each community (Burgess et al., 2003). Studies on resistance and efficacy are ongoing.

Resistance is often branded as the proximate cause of treatment failure when head lice are not eradicated by a pediculicide. However, these treat-

ment failures are more commonly the result of:

- Misdiagnosis (no active infestation, or misidentification).
- Non-compliance (not following treatment protocol).
- New infestation (lice acquired after treatment).
- Lack of ovicidal (egg-killing) or residual properties of the product (Burgess et al., 2003; Frankowski & Bocchini, 2010).

Pollack et al. (2000) found the most common reason for a conclusion of “resistance” was actually misdiagnosis; therefore, it could not be effectively treated with a pediculicide. Of the 555 samples sent in and initially identified as head lice, only 57.5% were correctly diagnosed and confirmed as such by an entomologist. Family identifications were only 47% correct, and physicians had the worst identification rate at 11% correct. The health care provider should consider resistance *after* assuring oneself the above factors have not contributed to the treatment failure.

Nurses as Change Agents

Pediatric nurses working both in and out of schools can lead the charge to help schools design evidence-based policies that respect the privacy and confidentiality of students yet promote successful treatment of infestations. NASN, in its position statement regarding pediculosis (NASN, 2011), described that school nurses are the key health care professionals in the provision of education and anticipatory guidance in all aspects of the management of pediculosis in the school setting. Additionally, school nurses play an important roll in helping local pharmacists, health care providers, and community health districts in updating their knowledge of current state of pediculosis science. Providing education to reduce the stigma of lice, clarify myths, and provide accurate information about effective treatment options, as well as appropriate referrals to health care providers, is important nursing care. Table 2 gives an example of a parent-teaching tool, designed by the author using the principles of health literacy to effectively dispute the common myths of lice management. NASN provides free, professionally produced, evidence-based parent and nurse educational tools in their *Lice Lessons* program (visit <http://www.nasn.org> for more information).

Table 2.
Lice 101: Myths and Realities about Head Lice

Definitions	
Lice: More than one louse.	Nit: Eggs, dead or alive, of a louse
Louse: Small insect that lives on the scalp.	Parasite: Lives off another, in this case the blood of humans.
Pediculosis: Having an infestation of lice.	Infestation: Having an insect present, in this case, in your head.
Myths	Truths
Head lice are easy to get.	Lice are spread only mainly by head-to-head contact. They are much harder to get than a cold, flu, ear infection, pink eye, strep throat, food poisoning, or impetigo.
You can get lice from your dog, guinea pig, or other animal.	Lice are species-specific. You can only get human lice from another human. You cannot get another animal's lice.
You can get head lice from hats and helmets.	Rarely, but possible. Hairbrushes, pillows, and sheets are also uncommon modes of transmission.
School is a common place for lice transmission.	School is an unlikely source of transmission. Much more common are family members, overnight guests, and playmates who spent a large amount of time together.
Poor hygiene contributes to lice.	Hygiene makes absolutely no difference. You get lice by close personal head-to-head contact with someone else that has lice, not by being dirty.
Lice can jump or fly from one person to another.	Lice can only crawl. They can neither fly nor jump. They must crawl from one person to another.
Any nits left in the hair can cause lice to come back.	Any nits farther away than one quarter to one half on the hair shaft are ALREADY HATCHED and pose no risk to others.
Eggs or nits can fall out of the hair, hatch, and cause lice in another person.	Nits are cemented to the hair and very hard to remove. They cannot fall off. Newly hatched lice must find a head quickly or will die.
Lice can live a long time.	Lice live only 1 to 2 days off the head. Each louse only lives about 30 days on the head.
All members of a family should be treated if one person has lice.	Only the person with lice should be treated. Lice shampoos are INSECTICIDES and can be dangerous if used incorrectly or too frequently. Household members and close contacts should be checked, but only treat those who actually have lice. The house should NOT be sprayed with insecticide, nor used on clothing or other items.
Checking a classroom when one student has lice can prevent lice from spreading.	Classroom transmission is EXCEEDINGLY RARE and checking students is a waste of valuable teaching time. Checking family members and close playmates is much more appropriate.
Avoiding lice is important as they spread disease.	Head lice do not spread any known disease. They are annoying and irritating, but not dangerous.

Even in light of evidence to the contrary, the lay public, including teachers and school administrators, often remain unconvinced of the need to remove “no-nit” policies and will be unmotivated to do so on their own. Further, some experts believe the only hope for true success is to focus on the children. “Teach them to think, to evaluate conflicting bits of information and to form logical and rational conclusions, and to be compassionate and caring” (R. Pollack, personal communication, May 21, 2014) to help children develop a new reality about head lice. Armed with the evidence,

school nurses are the perfect change agent to promote policy improvement to match what the evidence shows about the transmission and treatment of pediculosis. Not only can nurses provide education with every interpersonal encounter, they can lobby their school board. Following the examples set by leaders in school districts such as Oakland Unified, the author did just that (see Figure 6). Because many health care professionals are unaware of the realities of lice and newer lice treatments regimens, school nurses, as well as pediatric nurses in general, are in the ideal position to lead the charge!

References

- American Nurses Association (ANA). (2001). *Code of ethics for nurse*. Retrieved from <http://www.nursingworld.org/MainMenuCategories/EthicsStandards/CodeofEthicsforNurses/Code-of-Ethics.pdf>
- American Academy of Pediatrics (AAP). (2012). *Red book: 2012 report of the Committee of Infectious Diseases*. Elk Grove Village, IL: Author.
- American School Health Association (ASHA). (2000). *National Task Force on Confidential Student Health Information. Guidelines for protecting confidential student health information*. Kent, OH: ASHA.
- Bergren, M. (2001). HIPAA hoopla: Privacy and security of identifiable health info-

Figure 6.
Real Life Application

Using the evidence presented here, the author's district now allows both nits and live lice, does not send home specific alert letters, informs and educates parents/guardians at the end of the school day, and most importantly, provides copious teaching to families and children. In the four years hence, there has been no increase in infestation incidence, and although there is the occasional upset parent, overall the community has accepted these policies.

Source: Pontius, 2011.

- mation. *Journal of School Nursing*, 17(6), 336-340.
- Burgess, I., Pollack, R., & Taplin, D. (2003). *Cutting through controversy: Special report on the treatment of head lice*. Englewood, CO: Postgraduate Institute for Medicine.
- Centers for Disease Control and Prevention (CDC). (2013a). *Head lice information for schools*. Retrieved from <http://www.cdc.gov/parasites/lice/head/index.html>
- Centers for Disease Control and Prevention. (2013b). *Head lice treatment*. Retrieved from <http://www.cdc.gov/parasites/lice/head/treatment.html#otc>
- Centers for Disease Control and Prevention (CDC). (2013c). *Head lice biology*. Retrieved from www.cdc.gov/parasites/lice/head/biology.html
- Centers for Disease Control and Prevention (CDC) Public Health Image Library. (2014). *Homepage*. Retrieved from www.cdc.gov/phil/home.asp
- Frankowski, B., & Bocchini, J. (2010). Clinical report – Head lice. *Pediatrics*, 126(2), 392-403.
- Gordon, S. (2007). Shared vulnerability: A theory of caring for children with persistent head lice. *The Journal of School Nursing*, 23(5), 283-292.
- Gordon, S. (2009). *Head lice management in school settings*. Orlando, FL: Florida Association of School Nurses.
- Hootman, J. (2002). Quality improvement projects related to pediculosis management. *The Journal of School Nursing*, 18(2), 80-86.
- Loussouarn, G., Rawadi, C., & Genain, G. (2005). Diversity of hair growth profiles. *International Journal of Dermatology*, 44(Suppl. 1), 6-9.
- Meinking, T., & Taplin, D. (2011). Infestations. In L. Schachner, & R. Hansen (Eds.), *Pediatric dermatology* (pp. 1525-1583). Philadelphia: Mosby Elsevier.
- Mumcuoglu, K.B., Barker, F.C., Burgess, I.F., Combescot-Lang, C., Dagleish, R.C., Larsen K.S., ... Taylan-Ozkan, A. (2007). International guidelines for effective control of head louse infestations. *Journal of Drugs in Dermatology*, 6(4), 409-414.
- National Association of School Nurses (NASN). (2011). *Position Statement: Pediculosis in the school settings*. Silver Spring, MD: Author. Retrieved from <http://www.nasn.org/Portals/0/positions/2011pspediculosis.pdf>
- Pollack, R. (2012). Louse and nit removing salons. *IDmy Bug*. Retrieved from <http://idmybug.tumblr.com/post/17920155632/lice-nit-removing-salons>
- Pollack, R., Kiszewski, A., & Spielman, A. (2000). Overdiagnosis and consequent mismanagement of head louse infestations in North America. *Pediatric Infectious Disease Journal*, 19(8), 689-693.
- Pontius, D. (2011). Hats off to success: Changing head lice policy. *NASN School Nurse*, 26(6), 356-362.
- Speare, R., Cahill, C., & Thomas, G. (2003). Head lice on pillows, and strategies to make a small risk even less. *International Journal of Dermatology*, 42(8), 626-629.
- Speare, R., Thomas, G., & Cahill, C. (2002). Head lice are not found on floors in primary school classrooms. *Australian & New Zealand Journal of Public Health*, 26(3), 207-208.
- Williams, L., Reichert, M., MacKenzie, W., Hightower, A., & Blake, P. (2001). Lice, nits and school policy. *Pediatrics*, 107(5), 1011-1015.
- Yoon, K., Previte, D., Hodgdon, H., Poole, B., Kwon, D., Abo El-Ghar, G., ... Clark, J. (2014). Knockdown resistance allele frequencies in North American head louse populations. *Journal of Medical Entomology*. 51(2), 450-457. doi:10.1603/ME1313

Instructions For Continuing Nursing Education Contact Hours

Demystifying Pediculosis: School Nurses Taking the Lead

Deadline for Submission:
October 31, 2016

PED 1407

To Obtain CNE Contact Hours

1. For those wishing to obtain CNE contact hours, you must read the article and complete the evaluation through ***Pediatric Nursing's Web site*** at www.pediatricnursing.net/ce
2. Evaluations must be completed **online** by the above deadline. Upon completion of the evaluation, your CNE certificate for **1.3** contact hour(s) will be mailed to you.

Fees – Subscriber: Free Regular: \$20

Goal

To provide an overview of head lice and demystify the common misperceptions surrounding it.

Objectives

1. Define *Pediculosis capitis*.
2. List the common myths about head lice and nits.
3. Explain the implications these myths have on school children and their families.
4. Discuss ways school nurses may educate their community and schools about lice and nits, and thus, discourage the enforcement of "no-nit" policies.

Statement of Disclosure: The author(s) reported no actual or potential conflict of interest in relation to this continuing nursing education activity.

The *Pediatric Nursing* Editorial Board members reported no actual or potential conflict of interest in relation to this continuing nursing education activity.

This independent study activity is provided by **Anthony J. Jannetti, Inc.** (AJJ).

Anthony J. Jannetti, Inc. is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation.

Anthony J. Jannetti, Inc. is a provider approved by the California Board of Registered Nursing, Provider Number, CEP 5387.

Licenses in the state of California must retain this certificate for four years after the CNE activity is completed.

This article was reviewed and formatted for contact hour credit by Rosemarie Marmion, MSN, RN-BC, NE-BC, Anthony J. Jannetti, Inc., Education Director; and Judy A. Rollins, PhD, RN, *Pediatric Nursing* Editor.