

INDIANA 2018–19 Required and Recommended School Immunizations



| | REQUIRED | | RECOMMENDED |
|---|--|--|---|
| 3-5 years old | <ul style="list-style-type: none"> • 3 Hepatitis B • 4 DTaP (Diphtheria, Tetanus & Pertussis) • 3 Polio | <ul style="list-style-type: none"> • 1 Varicella (Chickenpox) • 1 MMR (Measles Mumps & Rubella) | <ul style="list-style-type: none"> • Annual Influenza |
| K-4th grade | <ul style="list-style-type: none"> • 3 Hepatitis B • 5 DTaP • 4 Polio | <ul style="list-style-type: none"> • 2 MMR • 2 Varicella • 2 Hepatitis A | <ul style="list-style-type: none"> • Annual Influenza |
| 5th grade | <ul style="list-style-type: none"> • 3 Hepatitis B • 5 DTaP • 4 Polio | <ul style="list-style-type: none"> • 2 MMR • 2 Varicella | <ul style="list-style-type: none"> • Annual Influenza • 2 Hepatitis A |
| 6th grade | <ul style="list-style-type: none"> • 3 Hepatitis B • 5 DTaP • 4 Polio • 2 MMR | <ul style="list-style-type: none"> • 2 Varicella • 2 Hepatitis A • 1 MCV4 (Meningococcal) • 1 Tdap (Tetanus, Diphtheria & Pertussis) | <ul style="list-style-type: none"> • Annual Influenza • 2 HPV (Human Papillomavirus) |
| 7th-11th grade | <ul style="list-style-type: none"> • 3 Hepatitis B • 5 DTaP • 4 Polio • 2 MMR | <ul style="list-style-type: none"> • 2 Varicella • 1 MCV4 (Meningococcal) • 1 Tdap (Tetanus, Diphtheria & Pertussis) | <ul style="list-style-type: none"> • Annual Influenza • 2 Hepatitis A • 2/3 HPV |
| 12th grade | <ul style="list-style-type: none"> • 3 Hepatitis B • 5 DTaP • 4 Polio • 2 MMR | <ul style="list-style-type: none"> • 2 Varicella • 2 Hepatitis A • 2 MCV4 • 1 Tdap | <ul style="list-style-type: none"> • Annual Influenza • 2 MenB (Meningococcal) • 2/3 HPV |

Hep B: The minimum age for the 3rd dose of Hepatitis B is 24 weeks of age.

DTaP: 4 doses of DTaP/DTP/DT are acceptable if 4th dose was administered on or after child's 4th birthday.

Polio: *3 doses of Polio are acceptable for all grade levels if the third dose was given on or after the 4th birthday and at least 6 months after the previous dose with only one type of vaccine used (all OPV or all IPV).

*For students in grades K–8th grade, the final dose must be administered on or after the 4th birthday and be administered at least 6 months after the previous dose.

Varicella: Physician documentation of disease history, including month and year, is proof of immunity for children entering preschool through 10th grade. Parental report of disease history is acceptable for grades 11–12.

Tdap: There is no minimum interval from the last Td dose.

MCV4: Individuals who receive dose 1 after their 16th birthday only need 1 dose of MCV4.

Hep A: The minimum interval between 1st and 2nd dose is 6 calendar months. 2 doses of Hep A is required for grades K–4, 6 and 12.

*For grades 5, and 7–11, 2 doses of Hep A vaccine are recommended.



Number under vaccine denotes the number of cumulative doses needed.