

Unit	Standards	Major Topics/Concepts
Ratios and Proportions	7.RP.A.1 7.RP.A.2 7.RP.A.3	<p>Compute unit rates associated with ratios of fractions, including ratios of lengths, areas, and other quantities measured in like or different units. <i>For example, if a person walks $\frac{1}{2}$ mile in each $\frac{1}{4}$ hour, compute the unit rate as the complex fraction $\frac{1/2}{1/4}$ miles per hour, equivalently 2 miles per hour.</i></p> <p>Recognize and represent proportional relationships between quantities.</p> <ul style="list-style-type: none"> ✓ Decide whether two quantities are in a proportional relationship (e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin). ✓ Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. ✓ Represent proportional relationships by equations. <i>For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as $t = pn$.</i> ✓ Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points $(0, 0)$ and $(1, r)$ where r is the unit rate. <p>Use proportional relationships to solve multi-step ratio and percent problems. <i>Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error.</i></p>
Operations with Rational Numbers	7.NS.A.1 7.NS.A.2 7.NS.A.3	<p>Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram.</p> <ul style="list-style-type: none"> ✓ Describe situations in which opposite quantities combine to make 0. ✓ Understand $p + q$ as the number located a distance q from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts. ✓ Understand subtraction of rational numbers as adding the additive inverse, $p - q = p + (-q)$. Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts. ✓ Apply properties of operations as strategies to add and subtract rational numbers.

Unit	Standards	Major Topics/Concepts
		<p>Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers.</p> <ul style="list-style-type: none"> ✓ Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the Distributive Property, leading to products such as $(-1)(-1) = 1$ and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts. ✓ Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then $-(p/q) = (-p)/q = p/(-q)$. Interpret quotients of rational numbers by describing real-world contexts. ✓ Apply properties of operations as strategies to multiply and divide rational numbers. ✓ Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats. <p>Solve real-world and mathematical problems involving the four operations with rational numbers. (Computations with rational numbers extend the rules for manipulating fractions to complex fractions.)</p>
1st Cumulative Benchmark (covering all content to this point)		
Expressions and Equations	7.EE.A.1 7.EE.A.2 7.EE.B.3 7.EE.B.4	<p>Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with rational coefficients.</p> <p>Understand that rewriting an expression in different forms in a contextual problem can provide multiple ways of interpreting the problem and how the quantities in it are related. <i>For example, shoes are on sale at a 25% discount. How is the discounted price P related to the original cost C of the shoes? $C - 0.25C = P$. In other words, P is 75% of the original cost for $C - 0.25C$ can be written as $0.75C$.</i></p> <p>Solve multi-step real-world and mathematical problems posed with positive and negative rational numbers presented in any form (whole numbers, fractions, and decimals).</p> <ul style="list-style-type: none"> ✓ Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate. ✓ Assess the reasonableness of answers using mental computation and estimation strategies. <p>Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities.</p> <ul style="list-style-type: none"> ✓ Solve contextual problems leading to equations of the form $px + q = r$ and $p(x + q) = r$, where p, q, and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. <i>For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?</i>

Unit	Standards	Major Topics/Concepts
		<p>✓ Solve contextual problems leading to inequalities of the form $px + q > r$ or $px + q < r$, where p, q, and r are specific rational numbers. Graph the solution set of the inequality on a number line, and interpret it in the context of the problem. <i>For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.</i> (Note that inequalities using $>$, $<$, \leq, \geq are included in this standard.)</p>
<p>Construction and Deconstruction of Geometric Figures</p>	<p>7.G.A.1 7.G.A.2</p>	<p>Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale.</p> <p>Draw geometric shapes with given conditions. Focus on constructing triangles from three measures of angles or sides, noticing when the conditions determine a unique triangle, more than one triangle, or no triangle.</p>
<p>2nd Cumulative Benchmark (covering all content to this point)</p>		
<p>Angles, Area, and Volume</p>	<p>7.G.B.3 7.G.B.4 7.G.B.5</p>	<p>Know the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle.</p> <p>Know and use facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle in a figure.</p> <p>Solve real-world and mathematical problems involving area, volume, and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.</p>
<p>Statistics</p>	<p>7.SP.A.1 7.SP.A.2 7.SP.B.3 7.SP.B.4</p>	<p>Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.</p> <p>Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. <i>For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.</i></p> <p>Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. <i>For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team; on a dot plot or box plot, the separation between the two distributions of heights is noticeable.</i></p>

Unit	Standards	Major Topics/Concepts
		<p>Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. <i>For example, decide whether the words in a chapter of a 7th grade science book are generally longer than the words in a chapter of a 4th grade science book.</i></p>
<p>Probability</p>	<p>7.SP.C.5 7.SP.C.6 7.SP.C.7 7.SP.D.8</p>	<p>Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.</p> <p>Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. <i>For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times but probably not exactly 200 times.</i></p> <p>Develop a probability model, and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy.</p> <ul style="list-style-type: none"> ✓ Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. <i>For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected.</i> ✓ Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. <i>For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?</i> <p>Summarize numerical data sets in relation to their context.</p> <ul style="list-style-type: none"> ✓ Give quantitative measures of center (median and/or mean) and variability (range and/or interquartile range), and describe any overall pattern and any striking deviations from the overall pattern with reference to the context in which the data were gathered. ✓ Know and relate the choice of measures of center (median and/or mean) and variability (range and/or interquartile range) to the shape of the data distribution and the context in which the data were gathered.
<p>Final Comprehensive Benchmark (covering all content)</p>		