



# California School Employee Tuberculosis (TB) Risk Assessment Questionnaire



(for pre-K, K-12 schools and community college employees, volunteers and contractors)

- Use of this questionnaire is required by California Education Code sections 49406 and 87408.6, and Health and Safety Code sections 1597.055 and 121525-121555.^
- The purpose of this tool is to identify **adults** with infectious tuberculosis (TB) to prevent them from spreading disease.
- **Do not repeat testing** unless there are **new risk factors since the last negative test**.
- **Do not treat for latent TB infection (LTBI) until active TB disease has been excluded:**  
*For individuals with signs or symptoms of TB disease or abnormal chest x-ray consistent with TB disease, evaluate for active TB disease with a chest x-ray, symptom screen, and if indicated, sputum AFB smears, cultures and nucleic acid amplification testing. A negative tuberculin skin test (TST) or interferon gamma release assay (IGRA) does not rule out active TB disease.*

Name of Person Assessed for TB Risk Factors: \_\_\_\_\_

Assessment Date: \_\_\_\_\_

Date of Birth: \_\_\_\_\_

| History of Tuberculosis Disease or Infection (Check appropriate box below) |   |
|--|---|
| <input type="checkbox"/>   | <b>Yes</b><br><ul style="list-style-type: none"> <li>• If there is a <u>documented</u> history of positive TB test or TB disease, then a symptom review and chest x-ray (if none performed in the previous 6 months) should be performed at initial hire by a physician, physician assistant, or nurse practitioner. If the x-ray does not have evidence of TB, the person is no longer required to submit to a TB risk assessment or repeat chest x-rays.</li> </ul> |
| <input type="checkbox"/>   | <b>No</b> (Assess for Risk Factors for Tuberculosis using box below)  |

| TB testing is recommended if <u>any</u> of the 3 boxes below are checked        |  |
|---|--|
| <input type="checkbox"/>  | <b>One or more sign(s) or symptom(s) of TB disease</b><br><ul style="list-style-type: none"> <li>• TB symptoms include prolonged cough, coughing up blood, fever, night sweats, weight loss, or excessive fatigue.</li> </ul>  |
| <input type="checkbox"/>  | <b>Birth, travel, or residence</b> in a country with an elevated TB rate for at least 1 month<br><ul style="list-style-type: none"> <li>• Includes countries <u>other than</u> the United States, Canada, Australia, New Zealand, or Western and North European countries.</li> <li>• Interferon gamma release assay (IGRA) is preferred over tuberculin skin test (TST) for non-US-born persons.</li> </ul> |
| <input type="checkbox"/>  | <b>Close contact</b> to someone with infectious TB disease during lifetime   |
| Treat for LTBI if TB test result is positive and active TB disease is ruled out |  |

^The law requires that a health care provider administer this questionnaire. A health care provider, as defined for this purpose, is any organization, facility, institution or person licensed, certified or otherwise authorized or permitted by state law to deliver or furnish health services. A Certificate of Completion should be completed after screening is completed (page 3).