



Texas Minimum State Vaccine Requirements for Child-Care Facilities

This chart summarizes the vaccine requirements incorporated in **Title 25 Health Services, §§97.61-97.72 of the Texas Administrative Code (TAC)**. This chart is not intended as a substitute for consulting the TAC, which has other provisions and details. The Department of State Health Services is granted authority to set immunization requirements by the **Human Resources Code, Chapter 42**.

2017-2018 Immunization Requirements for Child-Care Facilities

[Download this Chart as a PDF](#)

Age at which child must have vaccines to be in compliance:	Minimum Number of Doses Required of Each Vaccine							
	DTaP	Polio	HepB	Hib	PCV	MMR	Varicella	HepA
0 through 2 months	None	None	None	None	None	None	None	None
By 3 months	1 Dose	1 Dose	1 Dose	1 Dose	1 Dose	None	None	None
By 5 months	2 Doses	2 Doses	2 Doses	2 Doses	2 Doses	None	None	None
By 7 months	3 Doses	2 Doses	2 Doses	2 Doses ¹	3 Doses ²	None	None	None
By 16 months	3 Doses	2 Doses	2 Doses	3 Doses ¹	4 Doses ²	1 Dose ³	1 Dose ³	None
By 19 months	4 Doses	3 Doses	3 Doses	3 Doses ¹	4 Doses ²	1 Dose ³	1 Dose ³	None
By 25 months	4 Doses	3 Doses	3 Doses	3 Doses ¹	4 Doses ²	1 Dose ³	1 Dose ³	1 Dose ³
By 43 months	4 Doses	3 Doses	3 Doses	3 Doses ¹	4 Doses ²	1 Dose ³	1 Dose ³	2 Doses ³

¹ A complete Hib series is two doses plus a booster dose on or after 12 months of age (three doses total). If a child receives the first dose of Hib vaccine at 12 - 14 months of age, only one additional dose is required (two doses total). Any child who has received a single dose of Hib vaccine on or after 15 - 59 months of age is in compliance with these specified vaccine requirements. Children 60 months of age and older are not required to receive Hib vaccine.

² If the PCV series is started when a child is seven months of age or older or the child is delinquent in the series, then all four doses may not be required. Please reference the information below to assist with compliance:

- For children seven through 11 months of age, two doses are required.
- For children 12 - 23 months of age: if three doses have been received prior to 12 months of age, then an additional dose is required (total of four doses) on or after 12 months of age. If one or two doses were received prior to 12 months of age, then a total of three doses are required with at least one dose on or after 12 months of age. If zero doses have been received, then two doses are required with both doses on or after 12 months of age.
- Children 24 months through 59 months meet the requirement if they have at least three doses with one dose on or after 12 months of age, or two doses with both doses on or after 12 months of age, or one dose on or after 24 months of age. Otherwise, one